

The clinical safety and possibility of immediate implant placement in aggressive periodontitis patients in maxillary esthetic zone: an empty systematic review

Nermin Mohammed Ahmed Yussif¹
Ahmed Reda Abdel Rahman²
Mona Darhous²

¹ Oral Medicine and Periodontology Department, Faculty of Dentistry, October University for Modern Sciences and Arts (MSA University), Egypt

² Diagnosis, Oral Medicine and Periodontology Department, Faculty of Dentistry, Cairo University, Egypt

Address for correspondence:

Nermin Mohammed Ahmed Yussif, MSC
Oral Medicine and Periodontology Department
October University for Modern Sciences and Arts
(MSA University), Egypt
E-mail: dr_nermin_yusuf@yahoo.com

Summary

Purpose. To determine the clinical possibility and success of immediate implantation in aggressive periodontitis patients.

Methods. In August 2017, Pubmed, Cochrane central database, web of sciences, Lilacs and hand search were searched with no restrictions for language, publication date or follow-up period. All types of prospective studies reporting the prognosis of immediate implant placement were included. Outcomes were implant survival rate, complications during placement and postoperatively and marginal bone loss around implants.

Results. After screening of the titles, abstracts and full text of the reported studies, none of those met the *a priori* inclusion criteria.

Conclusion. There was no sufficient evidence regarding the survival rate of immediately placed dental implants in aggressive periodontitis. Long-term prospective double armed clinical studies are recommended.

KEY WORDS: immediate implant placement; aggressive periodontitis; empty systematic review.

Introduction

Achieving an attractive ideal smile has turned as the main goal for most individuals, especially patients suffering from periodontal diseases. Aggressive periodontitis, descriptively is an inscrutable disease that manifested by indeterminable periods of periodontal tissue destruction. It mainly affects youth with distinguishable progressive exacerbation periods. There is no intimate relationship between the amount of infective deposits and the rate of bone destruction. It results in unethical smile, malocclusion and marked functional discrepancy (1, 2).

The inability to identify the main causative factors behind such disease could be the reason behind missing the ideal treatment protocol. Treatment is usually the interplay of different factors. As per microbial hypothesis, *P. gingivalis* and *Aggregatibacter actinomycetem comitans* were guaranteed as the initiators (1, 3).

Afterwards, the genetic background was used to describe the familial behavior of the disease (4). Due to the high cost effectiveness, the genetic treatment is completely excluded especially in developing countries.

Additionally, several researches aimed to clarify the reason behind the hindered body response. Reduced functions of the immune cells and production of massive amounts of catabolic cytokines were reported. Furthermore, uncontrollable release of massive amounts of matrix-metalloproteinases was detected. Host modulators were recommended to control the release of the destructive agents and limit the disease activity (5, 6).

When dealing with aggressive periodontitis patients, immediate implant placement exhibits an effective modality that abbreviates the surgical time frame. It also provides proper implant positioning to counteract the advanced bone resorption following extraction (7, 8).

In general, peri-implantitis is a predictable event when treating patients with history of periodontitis in comparison to delayed technique. Meticulous precautions should be performed pre-operatively in order to control post-operative complications. Chemical and mechanical control of periodontitis is an essential step to control periodontitis regardless of its type (2, 9-11).

In 2005 and 2007, Mengel et al. (9, 10, 12, 13) have reported the long term success rate (83.33%) of immediately placed implants in aggressive periodontitis patients along 10 years follow-up.

On the other hand, Swierket et al. (2012) (11) assessed the peri-implant mucositis and peri-implantitis rate in aggressive periodontitis patients. They found that the implant survival rate was insignificant.

Immediate placement usually gains popularity attributable to the technique simplicity and limited surgical time. According to published literature, several systematic reviews discussed the efficacy of delayed implant placement aggressive periodontitis. Currently, the literature did not address short or long term treatment outcomes regarding immediate placement. The current systematic review aims to evaluate the evidence supporting the survival rate of immediately placed implants in patients with aggressive periodontitis supporting or limiting its performance.

Methodology and results

The Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA guidelines) was followed during the search protocol (Figure 1).




Figure 1 - PRISMA 2009 Flow Chart.

Focused question

The systematic search was performed aiming to answer the focused question:

“In aggressive periodontitis patients, what is the survival rate of the immediately placed dental implants versus periodontally healthy individuals in the maxillary esthetic zone?”

PICO elements

Participants (P)

Patients with hopeless teeth seek for immediate implant placement.

Types of interventions and comparisons (I & C)

The intervention was determined as immediate implant placement in aggressive periodontitis patients. While the comparator was the immediate placement in periodontally healthy individuals. Studies reporting delayed implant placement were excluded.

Types of outcome measures (O)

- 1st outcome: Survival rate of the immediately placed implants

- 2nd outcome: Radiographic interproximal marginal peri-implant bone loss (radiographic assessment)

Search strategy

Electronic data bases (PubMed, Cochrane central databases, JBI data base of systematic reviews and implementation reports, EBCAI evidence based communication assessment and intervention, Web of sciences and Lilacs) were searched in August 2017. Hand search in Google scholar data base, specialized journal databases (journal of periodontology, periodontology 2000, clinical periodontology) and bibliography of full text articles was performed. No language restrictions were applied. The search yielded 56 articles in total.

Publication selection

Titles and abstracts of the relevant studies were reviewed based on the inclusion criteria. Screening process was performed. Checking the full text of 21 studies was performed to exclude the irrelevant articles.

Study design selection and inclusion criteria

The relevant studies were categorized as randomized clinical

trials, non-randomized clinical trials, prospective studies and case studies. Prospective double arms clinical studies (RCTs, non-RCTs, cohort, case-control studies) that assessed the survival rate of immediately placed dental implants in aggressive periodontitis patients were included. No limitations concerning the follow-up period, language or publication date. Owing to reach trustable consensus and due to methodological heterogeneity, Authors excluded case reports and retrospective studies.

Unit inclusion criteria

Patients eligible for the trial must comply with all of the following: 1) Patients 18 years and older; 2) periodontally healthy individuals and aggressive periodontitis patients; 3) Medically healthy; 4) Non-smoker; 5) Immediate Implant placement; 6) Totally dentulous or partially edentulous patients.

Exclusion criteria

Twenty one full text studies were excluded. The reason behind exclusion was mainly the lack of short or long term articles discussing the technique. Deficient information concerning the study design was also great limitation.

Data synthesis

Table 1 - Excluded studies and the reasons behind.

| No. | Study name | Reason for exclusion |
|-----|--------------------------------------|--|
| 1. | De Boever & De Boever (2006) (14) | Prospective single arm clinical study evaluated delayed implant placement in aggressive periodontitis |
| 2. | Elamrousy et al. (2013) (15) | Prospective single arm clinical study evaluated immediate placement and restoration in all types of periodontitis (including chronic & aggressive periodontitis) |
| 3. | Elamrousy et al. (2014) (16) | Prospective single arm clinical study evaluated immediate placement and restoration in all types of periodontitis (including chronic & aggressive periodontitis) |
| 4. | Elsharkawy and El-Menoufy, 2013 (17) | Prospective clinical study evaluated immediate placement in patients with chronic periodontitis <i>versus</i> those placed in periodontally healthy patients |
| 5. | Furhauser et al. (2016) (18) | Prospective single arm clinical study evaluated immediate placement in patients periodontally healthy patients |
| 6. | Kehl et al. (2011) (19) | Prospective clinical study evaluated delayed placement in patients with aggressive periodontitis <i>versus</i> those placed in chronic periodontitis patients |
| 7. | Koh et al. (2011) (20) | Prospective clinical study evaluated immediate placement in healthy patients by placing implants at the level of the marginal bone crest <i>versus</i> placement 1 mm submarginal |
| 8. | Kolerman et al. (2016) (21) | The study was single arm prospective study evaluating the concept of immediate implant placement and nonfunctional loading for use in the esthetically sensitive anterior maxilla in periodontally healthy individuals |
| 9. | Koticha et al. (2012) (22) | Prospective clinical study evaluated immediate placement in patients periodontally healthy patients |

Reviewing and screening process was performed by the help of 2 Authors (NY and A.R). Both Authors extracted and revised data from the relevant studies independently. No studies retained. Table 1 showed the excluded studies and the reasons behind.

Discussion and summary of evidence

Immediate implantation is a critical treatment modality particularly in an uncontrolled destructive disease as aggressive periodontitis. Shortening of treatment schedule, cost effectiveness and appropriate implant placement are the main reasons behind preferring immediate placement. The target of the current systematic review was to assess the survival rate of immediately placed dental implants in aggressive periodontitis patients. Implant survival rate and radiographic interproximal marginal peri-implant bone loss were the measured outcomes. Comprehensive search resulted in no studies met the inclusion criteria. There was no evidence standing with or against the probable treatment procedure using prospective either short or long studies.

Despite the fact that numerous systematic reviews have discussed the role of aggressive periodontitis as a risk factor

to be continued

continued from Table 1

| | | |
|-----|---|--|
| 10. | Li et al. (2017) (23) | Prospective single arm study aimed to evaluate the feasibility and medium-term outcomes of immediate implant and rehabilitation based on the All-on-4 concept in patients with advanced GAP via clinical and radiographic analyses |
| 11. | Malo & Nobre (2008) (24) | Prospective clinical study evaluated delayed placement in patients partially edentulous patients with soft bone |
| 12. | Mengel et al. (2001) (25) | Prospective clinical study evaluated delayed placement in patients with aggressive periodontitis <i>versus</i> those placed in chronic periodontitis patients |
| 13. | Mengel & de-Jacoby (2005) ^a (12) | Prospective clinical study evaluated delayed placement following bone augmentation in patients with periodontitis <i>versus</i> those placed in healthy individuals |
| 14. | Mengel & de-Jacoby (2005) ^a (13) | Prospective clinical study evaluated delayed placement in patients with periodontitis (2 groups; aggressive periodontitis group and chronic periodontitis group) <i>versus</i> those placed in healthy individuals |
| 15. | Mengel et al. (2007) ^a (9) | Prospective clinical study evaluated delayed placement in patients with aggressive periodontitis <i>versus</i> those placed in healthy individuals |
| 16. | Mengel et al. (2007) ^b (10) | Prospective clinical study evaluated delayed immediate placement (2-4 weeks) in patients with aggressive periodontitis <i>versus</i> those placed in periodontally healthy patients |
| 17. | Mishra et al. (2011) (26) | Clinical report describes the delayed implant placement with aggressive periodontitis patients |
| 18. | Roccuzzo et al. 2010 (27) | Prospective clinical study evaluated delayed placement in patients with periodontally affected patients <i>versus</i> those placed in periodontally healthy patients |
| 19. | Schwartz-Arad et al. (2007) (28) | Prospective clinical study evaluated immediate placement in general |
| 20. | Slagter et al. (2016) (29) | Prospective clinical study evaluated immediate versus delayed placement with delayed loading in patients with labial bony defect |
| 21. | Swierkot et al. (2012) (11) | Prospective clinical study evaluated delayed placement in patients with aggressive periodontitis <i>versus</i> those placed in healthy individuals |

during delayed implantation, none evaluated the immediate placement. The start was in 2008, Al-Zahrani reported that the survival rate of dental implants placed in aggressive periodontitis patients has questionable survival rate. The results were owing to the impaired host response leading to uncontrolled progression of periodontitis. Regarding the included studies, 89-100% success rate resulted (30). While in 2012, Kim & Sung attributed the increased survival rate in delayed implants placed to adequate control of the infection and personalized maintenance program. The short-term success rate recorded 97.4%-100% while the long-term success recorded 83.3-96% success (31).

In 2014, a systematic review conducted assessing if aggressive periodontitis is considered a risk factor for implant failure. Nearly equal implant failure results were reported in aggressive periodontitis (3.97), healthy individuals and patients

with chronic periodontitis (4.0) (32). While in 2017, Theodoridis et al. (33) concluded that the survival rate of delayed placement in aggressive periodontitis patients were about 97.8% in comparison with periodontally healthy patients (100%). They also reported that the higher marginal bone loss in aggressive over chronic periodontitis and healthy individuals.

In the light of literature, confidence increased in placing delayed implants in aggressive periodontitis patients with better control of the infection and oral hygiene. In our opinion, the lack of immediate placement short or long term studies is the fear of the high failure rate.

Conclusion and implication for practice and research

Additional randomized clinical trials are needed to support

the clinical evidence of immediate implant placement in aggressive periodontitis patients with short, medium and long term outcomes.

Conflict of interest and source of funding statement

The Authors declare that they have no possible and the review was self-funded.

References

- Jain N, Jain G, Javed S, Iqbal Z, Talegaonkar S, Ahmad F, et al. Recent approaches for the treatment of periodontitis. *Drug Discover Today*. 2008;13:932-943.
- Pjetursson B, Helbling C, Weber H, Matulienė G, Salvi G, Brägger U, et al. Peri-implantitis susceptibility as it relates to periodontal therapy and supportive care. *Clin Oral Implants Res*. 2012;23:7:888-894.
- Slots J. Selection of antimicrobial agents in periodontal therapy. *J Periodont Res*. 2002;37:389-398.
- Nibali L, Donos N, Henderson B. Periodontal infectogenomics. *Journal of Medical Microbiology*. 2009;58:1269-1274.
- Suresh S. A new paradigm in Autoimmunity - role in periodontal disease. *Ind J Dent Advanc*. 2011;3:583-586.
- Bali D, Pandit N, Kathuria R, Bali A. Genetics and Aggressive Periodontal Disease: An Update Review. *J Oral Health Comm Dent*. 2012;6:2:97-101.
- Botticelli D, Renzi A, Lindhe J, et al. Implants in fresh extraction sockets: a pro-spective 5-year follow up clinical study. *Clin Oral Implants Res*. 2008;19:1226-1232.
- Lai Y, Kao S, Yeung T, Lee S. Rapid implant therapies: immediate placement and immediate restoration implant. *J Dent Sci*. 2009;4:1-6.
- Mengel R, Behle M, Flores-de-Jacoby L. Osseointegrated implants in subjects treated for generalized aggressive periodontitis: 10-year results of a prospective, long-term cohort study. *J Periodontol*. 2007;78:2229-2237(a).
- Mengel R, Kreuzer G, Lehmann K, Flores-de-Jacoby L. A telescopic crown concept for the restoration of partially edentulous patients with aggressive generalized periodontitis: a 3-year prospective longitudinal study. *Int J Periodontics Restorative Dent*. 2007;27:3: 231-239(b).
- Swierkot K, Lottholz P, Flores-de-Jacoby L, Mengel R. Mucositis, peri-implantitis, implant success, and survival of implants in patients with treated generalized aggressive periodontitis: 3- to 16-year results of a prospective long-term cohort study. *J Periodontol*. 2012;83:10:1213-1225.
- Mengel R, Flores-de-Jacoby L. Implants in patients treated for generalized aggressive and chronic periodontitis: a 3-year prospective longitudinal study. *J Periodontol*. 2005;76:4: 534-543(a).
- Mengel R, Flores-de-Jacoby L. Implants in regenerated bone in patients treated for generalized aggressive periodontitis: A prospective longitudinal study. *Int J Periodontics Restorative Dent*. 2005;25:331-341(b).
- De Boever A, De Boever J. Early colonization of nonsubmerged dental implants in patients with a history of advanced aggressive periodontitis. *Clin Oral Impl Res*. 2006;17:8-17.
- Elamrousy W, Nassar M, Ragheb A, Alnomany F, Marzok M. Radiographic Bone Changes around Immediately Placed Immediately Restored Dental Implants in Periodontally Compromised Sites. *Dentistry*. 2013;3:161.
- Elamrousy W, Nassar M, Alnomany F, Ragheb A, Markok R. Radiographic bone Changes around immediately placed immediate restored dental implants in periodontally compromised sites treated with Duo-Teck membrane. *The Journal of applied Sciences Research*. 2014;1:2, 85-96.
- Elsarkawy R, El-Menoufy H. Immediately Placed Implants in Periodontally Compromised Patients: A Prospective Clinical Study. *J Am Sci*. 2013;9:3:426-434.
- Furhauser R, Mailath-Pokorny G, Haas R, Busenlechner D, Watzek G, Pommer B. Immediate Restoration of Immediate Implants in the Esthetic Zone of the Maxilla Via the Copy-Abutment Technique: 5-Year Follow-Up of Pink Esthetic Scores. *Clinical Implant Dentistry and Related Research*. 2017;19:21.
- Kehl M, Swierkot K, Mengel R. Three-dimensional measurement of bone loss at implants in patients with periodontal disease. *J Periodontol*. 2011;82:689-699.
- Koh R, Oh T, Rudek I, Neiva G, Misch C, Rothman E, Wang H. Hard and Soft Tissue Changes After Crestal and Subcrestal Immediate Implant Placement. *J Periodontol*. 2011;82:1112-1120.
- Kolerman R, Mijiritsky E, Barnea E, Dabaja A, Nissan J, Tal H. Esthetic Assessment of Implants Placed into Fresh Extraction Sockets for Single-Tooth Replacements Using a Flapless Approach. *Clinical Implant Dentistry and Related Research*. 2016;Volume 00, Number 00.
- Koticha T, Fu J, Chan H, Wang H. Influence of Thread Design on Implant Positioning in Immediate Implant Placement. *J Periodontol*. 2012;83:1420-1424.
- Li S, Di P, Zhang Y, Lin Y. Immediate implant and rehabilitation based on All-on-4 concept in patients with generalized aggressive periodontitis: A medium-term prospective study. *Clin Implant Dent Relat Res*. 2017:1-13.
- Malo P, Nobre A. Flap vs. flapless surgical techniques at immediate implant function in predominantly soft bone for rehabilitation of partial edentulism: a prospective cohort study with a follow-up of 1 year. *Eur J Oral Implantol*. 2008;1:4:293-304.
- Mengel R, Schröder T, Flores-de-Jacoby L. Osseointegrated implants in patients treated for generalized chronic periodontitis and generalized aggressive periodontitis: 3- and 5-year results of a prospective long-term study. *J Periodontol*. 2001;72: 977-989.
- Mishra P, Mohamed J, Chandrasekaran S. Implants in periodontally compromised sites. *International Journal of Dental Clinics*. 2011;3:1: 100-101.
- Rocuzzo M, De Angelis N, Bonino L, Aglietta M. Ten year results of a three-arm prospective cohort study on implants in periodontally compromised patients. Part 1: Implant loss and radiographic bone loss. *Clin Oral Implants Res*. 2010;21:490-496.
- Schwartz-Arad D, Laviv A, Levin L. Survival of Immediately Provisionalized Dental Implants Placed Immediately Into Fresh Extraction Sockets. *J Periodontol*. 2007;78:219-223.
- Slagter K, den Hartog L, Bakker N, Vissink A, Meijer H, Raghoobar G. Immediate placement of dental implants in the esthetic zone: a systematic review and pooled analysis. *J Periodontol*. 2014;85:7:e241-250.
- Al-Zahrani M. Implant therapy in aggressive periodontitis patients: A systematic review and clinical implications. *Quintessence Int*. 2008;39:211-215.
- Kim K, Sung H. Outcomes of dental implant treatment in patients with generalized aggressive periodontitis: a systematic review. *The Journal of Advanced Prosthodontics*. 2012;4:4: 210-217.
- Monje A, Alcoforado G, Padial-Molina M, Suarez F, Lin G, Wang H. Generalized aggressive periodontitis as a risk factor for dental implant failure: a systematic review and meta-analysis. *J Periodontol*. 2014;85:10: 1398-1407.
- Theodoridis C, Grigoriadis A, Menexes G, Vouros I. Outcomes of implant therapy in patients with a history of aggressive periodontitis: A systematic review and meta-analysis. *Clin Oral Invest*. 2017;21:485-503.